



Western Washington University  
**Western CEDAR**

---

Salish Sea Ecosystem Conference

2020 Salish Sea Ecosystem Conference  
(Online)

---


Apr 21st, 9:00 AM - Apr 22nd, 4:45 PM

## **Squishy but not useless for energy balance: Energetic value of gelatinous zooplankton from the Salish Sea and adjacent waters**

Florian Lüskow

University of British Columbia, [flueskow@eoas.ubc.ca](mailto:flueskow@eoas.ubc.ca)

Follow this and additional works at: <https://cedar.wvu.edu/ssec>

 Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Natural Resources and Conservation Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

---

Lüskow, Florian, "Squishy but not useless for energy balance: Energetic value of gelatinous zooplankton from the Salish Sea and adjacent waters" (2020). *Salish Sea Ecosystem Conference*. 87.  
<https://cedar.wvu.edu/ssec/2020ssec/allsessions/87>

This Event is brought to you for free and open access by the Conferences and Events at Western CEDAR. It has been accepted for inclusion in Salish Sea Ecosystem Conference by an authorized administrator of Western CEDAR. For more information, please contact [westerncedar@wvu.edu](mailto:westerncedar@wvu.edu).



# Squishy but not useless for energy balance: Energetic value of gelatinous zooplankton from the Salish Sea and adjacent waters

Florian Lüsrow<sup>1,2</sup>, Moira D. Galbraith<sup>3</sup>, Brian P.V. Hunt<sup>1,2,4</sup>, R. Ian Perry<sup>5</sup>, Evgeny A. Pakhomov<sup>1,2,4</sup>

<sup>1</sup>Department of Earth, Ocean and Atmospheric Sciences, University of British Columbia, Vancouver, Canada   <sup>2</sup>Institute for the Oceans and Fisheries, University of British Columbia, Vancouver, Canada   <sup>3</sup>Institute of Ocean Sciences, Fisheries and Oceans Canada, Sidney, Canada   <sup>4</sup>Hakai Institute, Heriot Bay, Canada   <sup>5</sup>Pacific Biological Station, Fisheries and Oceans Canada, Nanaimo, Canada

E-mail: flueskow@eoas.ubc.ca, ORCID: 0000-0002-2100-7012, Poster-ID: 3351



## Introduction

- Gelatinous and soft-bodied zooplankton (GZ) are traditionally considered as ‘trophic dead ends’ (Hansson & Norman 1995)
- Due to their low energy contents per wet weight, GZ have been named ‘green tea of the sea’ (Hamilton 2016)
- There is growing awareness of importance of GZ in marine food webs (Hays *et al.* 2018) and for various other non-energetic purposes (Thiebot & McInnes 2020)
- Database listing elemental composition, organic and energy contents of various taxa is scattered, incomplete, and superficial in many cases
- We aim at the establishment of a comprehensive region-wide database compiling these data for a wide range of GZ species to inform energy-based food web models

## Methods

- For best describing the nutritional value of food items, a combination of several proxies is recommended (Chen & Li 2017; Machovsky-Capuska & Raubenheimer 2020)
- More than **1000 specimens** from **33 GZ species** were collected on **10 cruises** between **2014 and 2019** in the Salish Sea and adjacent waters (Figs. 1 and 2)
- Samples have been collected with Bongo and dip nets, Multinet, and pelagic trawl (Fig. 3)
- Specimens were sized and frozen on board and freeze-dried in the laboratory
- **Elemental composition (C, N)** was determined in elemental analyser
- **Organic content** measurement is based on dry matter (500 °C, 24 h)
- **Energy content** was estimated via published conversion factors (Platt *et al.* 1969; Schneider 1988) and based on bomb-calorimetry




Fig. 2




Fig. 5




Fig. 1




Fig. 7

### References

Chen L, Li C (2017) Different tolerances of jellyfish ephyrae (*Aurelia* sp.) and fish larvae (*Paralichthys olivaceus*) to nutrient limitations. *Mar Ecol Prog Ser* 569:1–13  
Hansson LJ, Norman B (1995) Release of dissolved organic carbon (DOC) by the scyphozoan jellyfish *Aurelia aurita* and its potential influence on the production of planktic bacteria. *Mar Biol* 121:527–532  
Hays GC, Doyle TK, Houghton JDR (2018) A paradigm shift in the trophic importance of jellyfish? *Trends Ecol Evol* 33(11):874–884  
Machovsky-Capuska GE, Raubenheimer D (2020) The nutritional ecology of marine apex predators. *Ann Rev Mar Sci* 12:361–387  
Platt T, Brown VM, Irwin B (1989) Caloric and carbon equivalents of zooplankton biomass. *J Fish Res Board Canada* 26(8):2345–2349  
Schneider G (1988) Chemische Zusammensetzung und Biomasseparameter der Ohrenqualle *Aurelia aurita*. *Helgol Meeresunters* 42:319–327  
Thiebot J-B, McInnes JC (2020) Why do marine endotherms eat gelatinous prey? *ICES J Mar Sci* 77(1): 58–71

### Acknowledgements

Captains and crews of CCGS J.P. Tully, CCGS W.E. Ricker, F/V Sea Crest, and R/V Professor Kaganovsky  
Stephen Healy, Sander Holthuisen, Henry Lam, Maureen Soon, and Lian E. Kwong for assistance with laboratory work

## Results

- Only a **set of net types with different mesh sizes** allows for huge species sampling variety in near and offshore areas in the present study
- **Marked and partly statistically significant differences in organic content and C/N ratio** among classes (Figs. 4 and 5) were identified
- Both depend highly on **sampled size range** (Fig. 6, e.g. *Aequorea* sp.) and **development stage composition** (Fig. 7, e.g. *Salpa aspera*)
- **Energy contents** vary between **< 0.1 and 22.8 kJ g DW<sup>-1</sup>** depended on class and method used (Fig. 8)
- Energy contents resulting from **bomb-calorimetry** are always low compared with conversion-used values
- **Organic content/energy content** pattern (based on Platt *et al.* 1969) highly similar to other GZ, but lower than values published for crustaceans (Fig. 9)

## Discussion

- ‘**Hidden diversity**’ in GZ in terms of organic and energy contents and elemental composition, which needs to be considered when used in food web models
- Size- and development stage dependency are considerable
- **Bomb-calorimetry** failed to confirm energy contents based on conversion factors; questioning either technique or validity of used conversions
- **Species-level diversity** is even bigger, but not shown here for clarity reasons
- Even if GZ have low energy content compared to crustaceans, **GZ and taxonomic variety** should not be neglected in nutritional studies

## Future Research

- Continuation of sample analyses from **cruises in 2020** is planned
- **Seasonality, various tissues, parasites, and further life cycle stages** are not considered yet
- Analysis of **phosphorus, vitamin, and micronutrient** contents is recommended
- Other gelatinous species of e.g. **siphonophores, polychaetes, and radiolarians** need to be analysed

